



# Global chemistry-climate modelling with EMAC

R. Deckert, R. Sausen, P. Jöckel, V. Aquila, S. Brinkop, U. Burkhardt, I. Cionni, M. Dall'Amico, M. Dameris, S. Dietmüller, V. Eyring, K. Gottschaldt, V. Grewe, J. Hendricks, M. Ponater, and M. Righi

Deutsches Zentrum für Luft- und Raumfahrt  
Institut für Physik der Atmosphäre  
Oberpfaffenhofen


High Performance Computing in Science and Engineering  
Garching, 8 - 9 December 2009



Deutsches Zentrum  
für Luft- und Raumfahrt e.V.  
in der Helmholtz-Gemeinschaft

# Introduction

## Global climate models



adapted from NOAA




Deutsches Zentrum  
für Luft- und Raumfahrt e.V.  
in der Helmholtz-Gemeinschaft

LRZ Garching 2009

# Introduction

## Global climate models: climate projections




source: IPCC



# Introduction

## Global climate models: climate projections



source: IPCC

# Introduction


Global climate models: no atmospheric chemistry

# Introduction

Global climate models: no atmospheric chemistry

- Future ozone layer?

21<sup>st</sup> November 2009



source: NASA



Deutsches Zentrum  
für Luft- und Raumfahrt e.V.  
in der Helmholtz-Gemeinschaft




# Introduction

Global climate models: no atmospheric chemistry

- Future ozone layer?
- Chemically triggered climate modifications?

21<sup>st</sup> November 2009



source: NASA



Deutsches Zentrum  
für Luft- und Raumfahrt e.V.  
in der Helmholtz-Gemeinschaft

# Introduction

Global chemistry-climate models (CCMs)

- Atmosphere ~~↔~~ deep ocean






# Introduction

## Global chemistry-climate models (CCMs)

- Atmosphere ~~↔~~ deep ocean
- Atmospheric physics ↔ atmospheric chemistry




# Introduction

Global chemistry-climate models (CCMs)

- Atmosphere ~~↔~~ deep ocean
- Atmospheric physics ↔ atmospheric chemistry

We use and refine a CCM



# Outline

- Model description
- Performance on LRZ/ALTIX
- Modelling activities on LRZ/ALTIX

# Description

CCM system **EMAC** (**E**CHAM/**M**ESSy **A**tmospheric **C**hemistry)

- Overall system by MPI for Chemistry, Mainz
- Fluid dynamics by MPI for Meteorology, Hamburg

# Description

CCM system **EMAC** (**E**CHAM/**M**ESSy **A**tmospheric **C**hemistry)

- Overall system by MPI for Chemistry, Mainz
- Fluid dynamics by MPI for Meteorology, Hamburg

EMAC follows standards → easily portable

- Fortran95 (ISO/IEC-1539-1)
- Message Passing Interface (MPI-2)
- NetCDF data format

# Description

CCM system **EMAC** (**E**CHAM/**M**ESSy **A**tmospheric **C**hemistry)

- Overall system by MPI for Chemistry, Mainz
- Fluid dynamics by MPI for Meteorology, Hamburg

EMAC follows standards → easily portable

- Fortran95 (ISO/IEC-1539-1)
- Message Passing Interface (MPI-2)
- NetCDF data format


EMAC is flexible

- Selectable spatial resolution
- Configurable chemistry scheme
- Modular structure (see <http://www.messy-interface.org>)

# Description

## Atmospheric primitive equations – spectral representation

- Spectral transformation (horizontal)
- Finite differences (vertical)
- Semi-implicit leap frog (time)






# Description

## Atmospheric primitive equations – spectral representation

- Spectral transformation (horizontal)
- Finite differences (vertical)
- Semi-implicit leap frog (time)

## Stiff system of ordinary differential equations – gaussian grid

Rosenbrock sparse-matrix technique with adaptive time step




# Description

## Atmospheric primitive equations – spectral representation

- Spectral transformation (horizontal)
- Finite differences (vertical)
- Semi-implicit leap frog (time)

## Stiff system of ordinary differential equations – gaussian grid

Rosenbrock sparse-matrix technique with adaptive time step



## Radiative transfer, tracer transport, other sub-models – gaussian grid

Miscellaneous techniques



# Description

## Diverse demands on hardware architecture ...

- Various numerical techniques

# Description

## Diverse demands on hardware architecture ...

- Various numerical techniques
- Distributed-memory domain decomposition
  - Spectral representation: spherical
  - Gaussian grid: latitude-longitude, ping-pong blocks
  - Tracer advection: vertical and latitude-longitude

# Description

## Diverse demands on hardware architecture ...

- Various numerical techniques
- Distributed-memory domain decomposition
  - Spectral representation: spherical
  - Gaussian grid: latitude-longitude, ping-pong blocks
  - Tracer advection: vertical and latitude-longitude
- Vectorisation of latitude-longitude blocks
  - To exploit cache sizes

# Description

## Diverse demands on hardware architecture ...

- Various numerical techniques
- Distributed-memory domain decomposition
  - Spectral representation: spherical
  - Gaussian grid: latitude-longitude, ping-pong blocks
  - Tracer advection: vertical and latitude-longitude
- Vectorisation of latitude-longitude blocks
  - To exploit cache sizes

→ EMAC valuable for benchmarking

# Performance on LRZ/ALTIX

Good performance, but gain limited

EMAC with  
detailed  
physics/chemistry

<i>vertical layers</i>	<i>CPUs</i>	<i>CPU time [kh/model year]</i>	<i>wall time [days/model year]</i>
<b>L41</b>	<b>64</b>	<b>5.4</b>	<b>3.5</b>
<b>L41</b>	<b>128</b>	<b>7.9</b>	<b>2.6</b>
<b>L90</b>	<b>64</b>	<b>12.0</b>	<b>7.9</b>
<b>L90</b>	<b>128</b>	<b>12.1</b>	<b>4.0</b>
<b>L90</b>	<b>256</b>	<b>poor</b>	<b>poor</b>





# Performance on LRZ/ALTIX

Good performance, but gain limited

- Communication between nodes (hardware specific)

EMAC with  
detailed  
physics/chemistry

<i>vertical layers</i>	<i>CPUs</i>	<i>CPU time [kh/model year]</i>	<i>wall time [days/model year]</i>
<b>L41</b>	<b>64</b>	<b>5.4</b>	<b>3.5</b>
<b>L41</b>	<b>128</b>	<b>7.9</b>	<b>2.6</b>
<b>L90</b>	<b>64</b>	<b>12.0</b>	<b>7.9</b>
<b>L90</b>	<b>128</b>	<b>12.1</b>	<b>4.0</b>
<b>L90</b>	<b>256</b>	<b>poor</b>	<b>poor</b>

# Performance on LRZ/ALTIX

Good performance, but gain limited

- Communication between nodes (hardware specific)
- Load imbalance from photo-chemistry (being tackled)

EMAC with  
detailed  
physics/chemistry

<i>vertical layers</i>	<i>CPUs</i>	<i>CPU time [kh/model year]</i>	<i>wall time [days/model year]</i>
<b>L41</b>	<b>64</b>	<b>5.4</b>	<b>3.5</b>
<b>L41</b>	<b>128</b>	<b>7.9</b>	<b>2.6</b>
<b>L90</b>	<b>64</b>	<b>12.0</b>	<b>7.9</b>
<b>L90</b>	<b>128</b>	<b>12.1</b>	<b>4.0</b>
<b>L90</b>	<b>256</b>	<b>poor</b>	<b>poor</b>

# Performance on LRZ/ALTIX

Good performance, but gain limited

- Communication between nodes (hardware specific)
- Load imbalance from photo-chemistry (being tackled)


L41 version: multi-decadal simulations sensible

EMAC with  
detailed  
physics/chemistry

<i>vertical layers</i>	<i>CPUs</i>	<i>CPU time [kh/model year]</i>	<i>wall time [days/model year]</i>
<b>L41</b>	<b>64</b>	<b>5.4</b>	<b>3.5</b>
<b>L41</b>	<b>128</b>	<b>7.9</b>	<b>2.6</b>
<b>L90</b>	<b>64</b>	<b>12.0</b>	<b>7.9</b>
<b>L90</b>	<b>128</b>	<b>12.1</b>	<b>4.0</b>
<b>L90</b>	<b>256</b>	<b>poor</b>	<b>poor</b>

# Development: new upper boundary


L90 expensive → our L41 version



# Development: new upper boundary

L90 expensive → our L41 version


- Appropriate for our purposes



# Development: new upper boundary

L90 expensive → our L41 version


- Appropriate for our purposes
- But: single layer for whole atmosphere above 25 km
- Impact on chemistry below



# Development: new upper boundary

Solution: highest layer with chemical parameterisation

- Calibration via L90 version






# Development: new upper boundary

Solution: highest layer with chemical parameterisation


- Calibration via L90 version



# Development: mixed-layer ocean

## Implementation of mixed-layer ocean

- Feedback: atmosphere ↔ mixed-layer ocean
- No feedback: mixed-layer ocean ↔ deep ocean



adapted from NASA



# Development: mixed-layer ocean


Mixed-layer ocean – additional computational demands

- Negligible for fix simulation length

# Development: mixed-layer ocean

Mixed-layer ocean – additional computational demands


- Negligible for fix simulation length
- However: spin-up about twenty years
- Much longer for deep ocean



# Development: mixed-layer ocean

Mixed-layer ocean – additional computational demands

- Negligible for fix simulation length
- However: spin-up about twenty years
- Much longer for deep ocean



# Development: Lagrangian formulation


Standard formulation of fluid dynamics: Eulerian

- Separate schemes for dynamics and tracer transport
- Tracer transport numerically diffusive

# Development: Lagrangian formulation

Standard formulation of fluid dynamics: Eulerian

- Separate schemes for dynamics and tracer transport
- Tracer transport numerically diffusive



Water vapour: ratio EMAC/satellite





# Development: Lagrangian formulation

Standard formulation of fluid dynamics: Eulerian

- Separate schemes for dynamics and tracer transport
- Tracer transport numerically diffusive

Implementation of full-Lagrangian tracer transport

- No numerical diffusion



Water vapour: ratio EMAC/satellite

# Development: Lagrangian formulation

Standard formulation of fluid dynamics: Eulerian

- Separate schemes for dynamics and tracer transport
- Tracer transport numerically diffusive

Implementation of full-Lagrangian tracer transport

- No numerical diffusion

**OPERATIONAL**



# Development: Lagrangian formulation

Standard formulation of fluid dynamics: Eulerian

- Separate schemes for dynamics and tracer transport
- Tracer transport numerically diffusive

Implementation of full-Lagrangian tracer transport

- No numerical diffusion

In development: full-Lagrangian dynamical core

- Finite-mass method (Gauger et al., 2000)
- Lagrangian air parcels with variable spatial extent
- Completely new approach in climate modelling

# Development: chemistry-transport mode

## Important question

- Impact of disturbed emission rates on atmospheric composition and climate
- E.g. human-made nitrogen oxide → near-surface ozone

# Development: chemistry-transport mode

## Important question

- Impact of disturbed emission rates on atmospheric composition and climate
- E.g. human-made nitrogen oxide → near-surface ozone

## Usually: difference of CCM simulations with un-/disturbed emission rates

- Noise from dynamical-chemical feedback
- Poor signal/noise ratio

# Development: chemistry-transport mode

## Important question

- Impact of disturbed emission rates on atmospheric composition and climate
- E.g. human-made nitrogen oxide → near-surface ozone

## Usually: difference of CCM simulations with un-/disturbed emission rates

- Noise from dynamical-chemical feedback
- Poor signal/noise ratio

## Our solution: switch off feedback

- Dynamics binary identical despite chemistry unequal
- Main challenge: polar stratospheric clouds

# Development: chemistry-transport mode

## Important question

- Impact of disturbed emission rates on atmospheric composition and climate
- E.g. human-made nitrogen oxide → near-surface ozone

Usually: difference of CCM simulation with and without emission rates

- Noise from dynamical-chemical feedback
- Poor signal/noise ratio

Our solution: switch on feedback

- Dynamical-chemical identical despite chemistry unequal
- Main challenge: polar stratospheric clouds

# Production simulations

Focus on model development:  
prerequisite for production simulations

→ Now feasible and going to start






# Production simulations

Climate impact of nitrogen oxide and sulfur dioxide from shipping


- Affects optical and microphysical cloud properties
- Mostly cooling



# Production simulations

## Aerosol aging

- Can result in ice nuclei
- Affects cirrus clouds



Number  
concentration of  
potential ice  
nuclei

# Summary

We use a chemistry-climate model (CCM)

- Diverse numerics / types of domain decomposition
- Good performance on LRZ/ALTIX
- But: gain limited due to communication between nodes

# Summary

We use a chemistry-climate model (CCM)

- Diverse numerics / types of domain decomposition
- Good performance on LRZ/ALTIX
- But: gain limited due to communication between nodes

Model development ...

- New upper boundary
- Mixed-layer ocean
- Lagrangian transport
- Chemistry-transport mode

# Summary

We use a chemistry-climate model (CCM)

- Diverse numerics / types of domain decomposition
- Good performance on LRZ/ALTIX
- But: gain limited due to communication between nodes

Model development ...

- New upper boundary
- Mixed-layer model
- Lagrangian transport
- Chemistry-transport mode

→ **Production going to start**